

Condition	Inclusion Criteria	Exclusion Criteria	Medicines	Leaflet
<b>Uncomplicated UTI in women</b>	16-64 years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> <li>• Urinary catheter</li> <li>• Recurrent (2x in last 6 months or 3x in last 12 months)</li> </ul>	<ul style="list-style-type: none"> <li>• Nitrofurantoin</li> </ul>	<a href="#">Target UTI leaflet</a>
<b>Shingles</b>	18+ years	<ul style="list-style-type: none"> <li>• Pregnant</li> </ul>	<ul style="list-style-type: none"> <li>• Aciclovir</li> <li>• Valaciclovir</li> </ul>	<a href="#">British Association of Dermatologists Shingles leaflet</a>
<b>Impetigo</b>	1+ years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> <li>• Bullous impetigo</li> <li>• Recurrent (2 or more in 12 months)</li> </ul>	<ul style="list-style-type: none"> <li>• Hydrogen peroxide cream</li> <li>• Fusidic acid cream</li> <li>• Flucloxacillin</li> <li>• Clarithromycin</li> <li>• Erythromycin</li> </ul>	<a href="#">British Association of Dermatologists Impetigo leaflet</a>
<b>Infected Insect Bites</b>	1+ years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> </ul>	<ul style="list-style-type: none"> <li>• Flucloxacillin</li> <li>• Clarithromycin</li> <li>• Erythromycin</li> </ul>	
<b>Acute Sore Throat</b>	5+ years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> </ul>	<ul style="list-style-type: none"> <li>• Penicillin V</li> <li>• Clarithromycin</li> <li>• Erythromycin</li> </ul>	<a href="#">Target RTI leaflet</a>
<b>Acute Sinusitis</b>	12+ years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> <li>• Immunosuppressed</li> <li>• Chronic sinusitis (&gt;12 weeks)</li> </ul>	<ul style="list-style-type: none"> <li>• Mometasone nasal spray</li> <li>• Fluticasone nasal spray</li> <li>• Penicillin V</li> <li>• Clarithromycin</li> <li>• Erythromycin</li> <li>• Doxycycline</li> </ul>	<a href="#">Target RTI leaflet</a>
<b>Acute Otitis Media</b>	1 – 17 years	<ul style="list-style-type: none"> <li>• Pregnant under 16</li> <li>• Recurrent (3x in 6 months or 4x in 12 months)</li> </ul>	<ul style="list-style-type: none"> <li>• Phenazone &amp; lidocaine ear drops</li> <li>• Amoxicillin</li> <li>• Clarithromycin</li> <li>• Erythromycin</li> </ul>	